

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

Modeling He-rich subdwarfs through the Hot-Flasher scenario: First Results

Miller Bertolami M. M.^{1,2} and Althaus L. G.^{1,2}

¹ *Facultad de Ciencias Astronómicas y Geofísicas, UNLP, Argentina*

² *Instituto de Astrofísica La Plata, UNLP-CONICET, Argentina*

Abstract.

We present first results from evolutionary simulations aimed at exploring the Hot-Flasher scenario for the formation of H-deficient subdwarf stars. The two types of late hot flashers that lead to He-enriched surfaces, “deep” and “shallow” mixing cases, are investigated for different metallicities.

Introduction: Hot subluminous stars are an important population of faint blue stars which can be roughly grouped into the cooler sdB stars and the hotter sdO stars. Subluminous O, B stars have been identified with stars populating the hot end of the horizontal branch of some globular and open clusters. While sdB stars form a nearly homogeneous spectroscopical class, a large variety of spectra is observed among the sdO stars (Stroeer et al. 2007). This variety of spectra, and the helium enhanced surface abundances pose a challenge to stellar evolution. As a consequence some non-canonical evolutionary scenarios have been proposed for their formation. Among them the “Hot Flasher scenario” (Castellani & Castellani 1993, D’Cruz et al. 1996, Sweigart 1997) offers one of the best possibilities (however, see Heber, this proceedings) for explaining their formation, while also explaining the correlation between the existence of blue hook stars and helium-core white swarfs (Hansen 2005, Kalirai et al. 2007).

Description of the work and Results: We performed full evolutionary simulations of the two flavors of the Hot-Flasher scenario, “deep” (DM) and “shallow” (SM) mixing cases (Lanz et al. 2004), for two different metallicities ($Z=0.001, 0.03$). Initial values were chosen to obtain ages, at the moment of the helium core flash, similar to those observed in globular clusters. Initial abundances were chosen in agreement with the evolution of galactic abundances presented by Flynn (2004). Surface abundances of our simulations are displayed in Table 1 and Fig. 1 shows the $\log T_{\text{eff}}-g$ values of the sequences.

We find that all our sequences display $C/N > 1$ (by mass), and thus should be considered “carbon rich” contrary to what is inferred at some He-sdO stars, which present both $C/N > 1$ and $C/N < 1$ surface abundances (Stroeer et al. 2007). Also, consistent with previous works, we find that the range of masses for late hot flashers is narrow and slightly dependent on mass. As noted by Lanz et al. (2004) we also find that the proportion of shallow mixing cases is dependent on mass, being $< 5\%$ at $Z = 0.001$ and $\sim 17\%$ for $Z=0.03$.

The surface temperatures and gravities of our sequences during the He-burning phase are cooler and show a narrow range of gravity values, which are not consistent with those inferred in He-SdO stars by Stroeer et al. (2007). Consequently our preliminary results do not seem to support the hot flasher

ZAMS: Mass= 0.88 $M_{\odot}$, X/Y/Z= 0.769/0.230/0.001. Age at Helium Flash= 12404 Myr

Final Mass [$M_{\odot}$]	H	He	^{12}C	^{13}C	N	O
0.4915 (SM)	0.21	0.323	0.038	1.5×10^{-6}	0.0001	0.0014
0.4910 (DM)	1.4×10^{-5}	0.955	0.0284	0.0076	0.0086	7×10^{-5}
0.4815 (DM)	1.2×10^{-6}	0.965	0.0139	0.0047	0.0163	4×10^{-5}

ZAMS: Mass= 1.04 $M_{\odot}$, X/Y/Z= 0.668/0.302/0.030. Age at Helium Flash= 12863 Myr

Final Mass [$M_{\odot}$]	H	He	^{12}C	^{13}C	N	O
0.4657 (SM)	0.512	0.457	0.00297	0.000121	0.00698	0.0103
0.4649 (SM)	0.026	0.905	0.0362	1.3×10^{-5}	0.00622	0.0019
0.4633 (SM)	0.00195	0.927	0.0371	0.00344	0.00694	0.00158
0.4628 (DM)	1.3×10^{-5}	0.933	0.0424	0.00155	0.00705	0.00154
0.4531 (DM)	4.5×10^{-6}	0.939	0.031	0.00474	0.01205	0.00198

Table 1. Surface abundances (by mass fractions) of the sequences calculated in this work. The choice of the initial values of the sequences (at ZAMS) are shown on top of each table.




Figure 1. Comparison of the values of T_{eff} and g of our sequences for different metallicities (right panel: $Z=0.03$, left panel: $Z=0.001$) as compared with the values inferred by Stroeger et al. (2007).

origin for these stars unless strong systematics are present in the observational inferences (Lanz et al. 2004, Heber private communication).

Acknowledgments. M3B thanks the organizers of the Third Conference in Hydrogen Deficient Stars for financial assistance that allowed him to participate in the conference.

References

- Brown, T., Sweigart, A., Lanz, T., Landsman, W., Hubeny, I. 2001, ApJ, 562, 368
- Castellani & Castellani 1993, A&A, 457, 569
- Cassisi, S., Schlattl, H., Salaris, M., Weiss, A. 2003, 582, L43
- D'Cruz, N., Dorman, B., Rood, R., O'Connell, R. 1996, ApJ, 466, 359
- Flynn, C. 2004, Publication of the Astron. Soc. of Australia, 21, 126
- Hansen, B. 2005, ApJ, 635, 522
- Kalirai, J., Bergeron, P., Hansen, B., Kelson, D., Reitzel, D., Rich, R., Harvey, R. 2007, arXiv:0705.0977v1 [astro-ph]
- Lanz, T., Brown, T., Sweigart, A., Hubeny, I., Landsman, W. 2004, ApJ, 602, 342
- Stroeger, A., Heber, U., Lisker, T., Napiwotzki, R., Dreizler, S., Christlieb, N., Reimers, D. 2007, A&A, 462, 269
- Sweigart, A. 1997, Third Conference on Faint Blue Stars, 3